

## RESEARCH ARTICLE

## Open Access



# Molybdate transporter ModABC is important for *Pseudomonas aeruginosa* chronic lung infection

Simone Périnet<sup>1,2</sup>, Julie Jeukens<sup>1,2</sup>, Irena Kukavica-Ibrulj<sup>1,2</sup>, Myriam M. Ouellet<sup>1,3,4</sup>, Steve J. Charette<sup>1,3,4</sup> and Roger C. Levesque<sup>1,2\*</sup>

## Abstract

**Background:** Mechanisms underlying the success of *Pseudomonas aeruginosa* in chronic lung infection among cystic fibrosis (CF) patients are poorly defined. The *modA* gene was previously linked to in vivo competitiveness of *P. aeruginosa* by a genetic screening in the rat lung. This gene encodes a subunit of transporter ModABC, which is responsible for extracellular uptake of molybdate. This compound is essential for molybdoenzymes, including nitrate reductases. Since anaerobic growth conditions are known to occur during CF chronic lung infection, inactivation of a molybdate transporter could inhibit proliferation through the inactivation of denitrification enzymes. Hence, we performed phenotypic characterization of a *modA* mutant strain obtained by signature-tagged mutagenesis (STM\_ *modA*) and assessed its virulence in vivo with two host models.

**Results:** The STM\_ *modA* mutant was in fact defective for anaerobic growth and unable to use nitrates in the growth medium for anaerobic respiration. Bacterial growth and nitrate usage were restored when the medium was supplemented with molybdate. Most significantly, the mutant strain showed reduced virulence compared to wild-type strain PAO1 according to a competitive index in the rat model of chronic lung infection and a predation assay with *Dictyostelium discoideum* amoebae. As the latter took place in aerobic conditions, the in vivo impact of the mutation in *modA* appears to extend beyond its effect on anaerobic growth.

**Conclusions:** These results support the *modABC*-encoded transporter as important for the pathogenesis of *P. aeruginosa*, and suggest that enzymatic machinery implicated in anaerobic growth during chronic lung infection in CF merits further investigation as a potential target for therapeutic intervention.

**Keywords:** ModA, Molybdate, *Pseudomonas aeruginosa*, Cystic fibrosis, Anaerobic conditions, Biofilm, Chronic lung infection, Animal model

## Background

*Pseudomonas aeruginosa* is an environmental bacterium and the most common cause of chronic lung infection among cystic fibrosis (CF) patients [1]. Its success in causing opportunistic infection and its persistence capacities are presumably attributable to a large 6.3–6.9 Mbp genome regulated by more than 550 transcriptional regulators, which allow adaptation to diverse environments

and growth conditions [2, 3]. *P. aeruginosa* also produces a wide repertoire of molecules and sensors for nutrient uptake, adhesion, mobility, biofilm formation and antibiotic resistance that are all key components of in vivo proliferation [4].

It was shown that the CF mucus is oxygen-depleted [5] and may carry strict anaerobes [6]. Hence, during chronic lung infection in CF, *P. aeruginosa* is exposed to microaerophilic [7] or anaerobic conditions [8], which are well suited for biofilm formation [9]. Denitrification, the main source of energy production under anaerobic conditions, is based upon the reduction of

\*Correspondence: [rclevesq@ibis.ulaval.ca](mailto:rclevesq@ibis.ulaval.ca)

<sup>1</sup> Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec, QC, Canada

Full list of author information is available at the end of the article

oxidized forms of nitrogen (preferentially nitrates,  $\text{NO}_3$ ) by metalloenzymes such as nitrate reductase [10, 11]. Molybdate ( $\text{MoO}_4^{2-}$ ) is the usable form of trace-element molybdenum (Mo) [12] and resembles sulfate, phosphate, tungstate and vanadate in molecular size, shape, charge and hydrogen-binding properties [13]. Mo is incorporated into the molybdenum cofactor (MoCo) and was found to be essential for the activity of molybdoenzymes. These enzymes catalyze various oxidation/reduction reactions and are implicated in the metabolism of nitrogen, carbon and sulfur. All nitrate reductases required for *P. aeruginosa* anaerobic growth require a MoCo cofactor [11], which can contain either Mo or tungsten (W) [14].

A previous PCR-based signature-tagged mutagenesis (PCR-STM) mutant screen allowed the identification of 148 genes presumably essential for in vivo survival of *P. aeruginosa* PAO1 in a rat model of chronic lung infection [15]. One of these genes was *modA*. In *E. coli*, molybdate and tungstate are internalized using an ATP-binding cassette transporter, ModABC [16], where ModA is the periplasmic binding protein with a high affinity for Mo/W, ModB is the integral membrane channel protein and ModC is the energizer protein [17]. In *P. aeruginosa*, *modA* was recently demonstrated to be essential for molybdate acquisition and anaerobic growth using a deletion mutant [18]. Here, with the goal of gaining further information on the role of ModABC in the virulence of *P. aeruginosa*, we present a characterization of the PAO1 transposon mutant STM\_ *modA*. We then assess the relevance of *modA* as a therapeutic target by testing virulence attenuation of STM\_ *modA* in two host models: the rat model of chronic lung infection, which best represents the context of CF lung infections, and the amoeba predation assay, which takes place in aerobic conditions. Results suggest that *modA* is important for the pathogenesis of *P. aeruginosa* in both host models.

## Methods

### Bacterial strains, plasmids, primers and culture conditions

Bacterial strains and plasmids used in this study are listed in Table 1. The STM\_ *modA* mutant was constructed and identified in the context of PCR-based STM using strain PAO1 [15, 19, 20], a laboratory strain previously shown to be appropriate and comparable to a CF isolate (LESB58) for the study of virulence using the rat model of chronic lung infection [21]. Briefly, a tagged-miniTn5Km2 suicide plasmid was transferred into strain PAO1 by conjugation, generating thousands of mutants, each containing a unique random mutation. Mutant strains were then negatively screened in pools of 72 in the rat model of chronic lung infection to identify defective mutants for in vivo maintenance via multiplex PCR. Mutation of *modA* in the STM\_ *modA* mutant strain was confirmed by cloning using a plasmid rescue based on the miniTn5Km2 antibiotic resistance marker followed by sequencing and alignment on the PAO1 genome available at the *Pseudomonas* genome database [22].

For routine cultures, *P. aeruginosa* and *E. coli* were grown aerobically at 37 °C in tryptic soy broth (TSB) or Luria–Bertani (LB) broth (EMD Serono). When needed, culture medium was supplemented with 1.5 (w/v) % bacto agar, kanamycin (Km 150  $\mu\text{g ml}^{-1}$  for STM\_ *modA*; Calbiochem), ampicillin (Ap 100  $\mu\text{g ml}^{-1}$  for DH10B transformants; Sigma-Aldrich) or carbenicillin (Cb 200  $\mu\text{g ml}^{-1}$  for *P. aeruginosa* transformants; Thermo-Fisher).

For growth experiments, *P. aeruginosa* was grown aerobically under vigorous shaking (250 rpm) in 100 ml LB using a 1:10 pre-culture to Erlenmeyer volume ratio. For anaerobic growth, nephelo flasks sealed with silicone stoppers containing 100 ml LB were flushed with argon for 45 min and needle-inoculated with 1 ml of pre-culture before incubation with weak agitation (120 rpm) to prevent precipitation. Media for aerobic and anaerobic growth were supplemented with potassium nitrate ( $\text{KNO}_3$ ; 15 mM; Merck) and, when needed, sodium

**Table 1** Strains and plasmids

Strain or plasmid	Relevant characteristics	Source or reference
<i>E. coli</i> strain		
NEB 5a	Chemically competent cells	New England Biolabs
<i>P. aeruginosa</i> strains		
PAO1	PAO1293, $\text{Cm}^S$ , wild-type, derivative of PAO2 which originates from PAO1	[48]
STM_ <i>modA</i>	PAO1 STM_ <i>modA</i> ::miniTn5-Km2 mutant inactivating <i>modA</i> (PA1863), $\text{Km}^R$	[15]
Complemented STM_ <i>modA</i>	PAO1 STM_ <i>modA</i> with pUCP19:: <i>modABC</i> plasmid (PA1863, PA1862, PA1861)	This study
Plasmids		
pUCP19	$\text{Cb}^R$ , cloning vector and used for CI	[49]
pUCP19:: <i>modABC</i>	$\text{Cb}^R$ , $\text{Km}^R$ , <i>modABC</i> genes on pUCP19 vector	This study

molybdate (100  $\mu$ M; Sigma-Aldrich). Growth was monitored by spectrophotometric measurements of optical density at 600 nm. Growth experiments were repeated three times.

Restriction enzymes, Q5 polymerase and Gibson assembly cloning kit were purchased from New England Biolabs. The QIAprep Spin Miniprep Kit (Qiagen) was used for plasmid isolation and the DNeasy Blood and Tissue kit (Qiagen) was used for genomic DNA isolation. PCR reactions were performed in an iCycler (Bio-Rad); primers used in this study are listed in Additional file 1.

#### Spectrophotometric determination of nitrate in culture medium after anaerobic growth

To determine if the STM\_ *modA* mutant was able to use nitrates during anaerobic growth, the concentration of residual nitrates in the medium was quantified after overnight growth based on the principle of chemical reduction of nitrate and its spectrophotometric detection using the Griess reaction [23]. Vanadium (III) chloride (Sigma-Aldrich) was used for the reduction of nitrates to nitrites. Sulfanilamide and N-(1-naphthyl)-ethylenediamine (Sigma-Aldrich) were used in the composition of the Griess reagent. Nitrite concentration was subtracted from the total nitrate and nitrite concentration to obtain the nitrate concentration alone.

#### Complementation of the STM\_ *modA* mutation

Cloning of the *modABC* operon into the pUCP19 vector was done using the Gibson Assembly Master Mix (New England Biolabs) following manufacturer's instructions. Two overlapping PCR fragments (Additional file 1) containing the entire *modABC* operon plus a 448 nt upstream region (2.1 kb final insert size) and an overlapping section of the *KpnI* and *PstI* restriction sites were ligated with the digested vector. The recombinant plasmid was introduced by heat shock into *E. coli* NEB 5 $\alpha$  chemically competent cells (Table 1). The plasmid transfer was first confirmed by *EcoRI* and *PstI* digestion followed by DNA sequencing. The plasmid was then recovered and electroporated into the *P. aeruginosa* STM\_ *modA* mutant. Plasmid insertion in the mutant strain was confirmed using digestions with the same restriction enzymes.

#### Biofilm formation assay

Multiple phenotypic tests were performed for this study (see reference [24] for more information on the protocols used), but only biofilm formation showed variation among strains. To measure the quantity of biofilm produced by wild-type strain PAO1, STM\_ *modA*, and the complemented STM\_ *modA*, a 96-plate rapid biofilm formation assay was performed as previously described [25]. Briefly, strains were grown overnight in LB supplemented

when needed with Km or Cb. The M63 culture medium for biofilm formation [25] was supplemented when needed with 100  $\mu$ M sodium molybdate and 15 mM KNO<sub>3</sub>. Biofilms were stained with crystal violet after a 6-hour static incubation time. Biofilm formation was quantified after stain dissolution in 2  $\times$  200  $\mu$ l 95 % v/v ethanol [26]. The experiment was done three times, with eight repetitions for each strain and condition. Statistical significance was assessed using an ANOVA in GraphPad Prism 6.0.

#### Ethics statement

The rat model in this study was used in a protocol approved by the "Comité de protection des animaux de l'Université Laval" (certificate 2011194-3, IACUC is a Canadian Council on Animal Care certificate holder).

#### Preparation of agarose bead-embedded bacteria and in vivo competitive index

The previously described rat model of chronic lung infection [27] using agarose bead-embedded bacteria was optimized as described elsewhere [20]. Sprague–Dawley rats of 450–500 g in weight were sedated (ketamine/xylazine IP injection, 10 mg/100 g) and a local anesthetic (lidocaine) was applied to the vocal cords. Animals were then intubated and inoculated with 120  $\mu$ l of bead preparation containing PAO1 + pUCP19 (Cb<sup>R</sup>) strain and the STM\_ *modA* mutant in equal parts ( $1.6 \times 10^6$  CFU ml<sup>-1</sup> per strain) for competitive index (CI) determination. At day 7 post-infection, rats were euthanized by barbiturate overdose (Euthanyl IP injection, 120 mg/kg) and homogenized lungs were diluted and plated in triplicate on TSA to quantify the total number of viable *P. aeruginosa* cells (TSA supplemented with 200  $\mu$ g Cb ml<sup>-1</sup> for the wild-type selection or TSA supplemented with 150  $\mu$ g Km ml<sup>-1</sup> for the mutant selection). In vivo CI was calculated as the ratio of mutant to wild-type bacteria recovered in vivo and adjusted according to the input ratio. The final CI data was represented as the geometric mean for each group of six animals and statistical significance was assessed with a two-tailed Mann–Whitney test on GraphPad Prism 6.0 software.

#### Amoeba predation assay

A predation assay was used to determine the bacterial capacity to resist to amoeba grazing [28]. The *D. discoideum* amoeba (DH1-10) was grown in HL5 medium (14.3 g l<sup>-1</sup> of bactopectone, 7.15 g l<sup>-1</sup> of yeast extract, 18 g l<sup>-1</sup> of D-(+)-monohydrate maltose, 0.641 g l<sup>-1</sup> of Na<sub>2</sub>HPO<sub>4</sub>•2H<sub>2</sub>O, and 0.490 g l<sup>-1</sup> of KH<sub>2</sub>PO<sub>4</sub>) [29] supplemented with 15  $\mu$ g ml<sup>-1</sup> tetracycline. The confluence of amoebae was about 60 % the day of the experiment. A volume of 300  $\mu$ L of *P. aeruginosa* liquid pre-culture

(OD<sub>600</sub> = 0.9) was spread in a uniform lawn on a Petri dish containing SM 1/10 agar (1 g l<sup>-1</sup> of bactopeptone, 0.1 g l<sup>-1</sup> of yeast extract, 0.22 g l<sup>-1</sup> of KH<sub>2</sub>PO<sub>4</sub>, 0.1 g l<sup>-1</sup> of K<sub>2</sub>HPO<sub>4</sub>, 0.1 g l<sup>-1</sup> of MgSO<sub>4</sub>·7H<sub>2</sub>O, 2 g l<sup>-1</sup> of Bacto agar, and 1 g l<sup>-1</sup> of glucose). Serial 1/10 dilutions of 50,000 amoeba cells/5 µL down to 0 cell/5 µL were deposited on the bacterial lawn. Petri dishes were incubated at 21 °C for 7 days and phagocytic plaques due to amoeba grazing were monitored. The experiment was repeated three times.

## Results and discussion


### Genetic characterization of the *P.aeruginosa* STM\_ *modA* mutant


Sequencing of the mutated *modA* gene showed that the miniTn5 insertion and its typical 9-pb flanking duplication [30] were inserted in the open reading frame (ORF) of *modA* at position 540 (Fig. 1), and introduced a stop codon assumed to prematurely interrupt translation of the protein. In the 6.3 Mbp genome of *P. aeruginosa* strain PAO1, *modA* (PA1863) is encoded in a putative operon with *modB* (PA1862) and *modC* (PA1861) (Fig. 1). Their products correspond to the 26.4 kDa (252 AA) molybdate-binding periplasmic protein precursor ModA, molybdenum membrane transport protein ModB (24.4 kDa) and ATPase ModC (39.8 kDa). These annotations are based on AA identity with ModABC from *E.coli* [31], where the *modABC* transcription unit codes for an ATP-binding cassette (ABC) transporter of molybdate, and were confirmed by Pederick and colleagues [18] using a *modA* deletion mutant. Since ABC transporters require a binding protein to present the substrate to the cargo membrane protein(s) [32], ModABC is expected to be non-functional in the STM\_ *modA* mutant.

### *modA* is required for anaerobic growth and nitrate utilization

Under aerobic conditions, growth of STM\_ *modA* was equivalent to that of wild-type PAO1. In contrast, under anaerobic conditions where NO<sub>3</sub> was provided as the terminal electron acceptor, the STM\_ *modA* mutant was unable to perform productive growth (Fig. 2a). However, when the growth medium was supplemented with 100 µM of sodium molybdate, mutant growth was restored to wild-type levels. Molybdate supplementation had no effect on PAO1 growth. Wild-type *modABC* was provided in *trans* to the mutant strain for complementation analysis using the expression vector pUCP19. The complemented STM\_ *modA* strain showed PAO1-like in vitro growth levels under anaerobic conditions (Fig. 2b), which confirmed that the disruption of *modA* is responsible for the STM\_ *modA* mutant phenotype in anaerobic conditions and that the ModABC transporter is essential for anaerobic growth. To confirm that this anaerobic growth defect was due to the malfunction of one or all nitrate reductases, residual nitrates in the growth medium were quantified (Fig. 2c). In the presence of nitrate reductase activity, nitrate concentration is expected to decrease due to bacterial uptake and subsequent reduction. After overnight growth, residual nitrates in the growth medium of wild-type PAO1 were completely consumed. In contrast, for the STM\_ *modA* mutant, nitrate concentration at the end of the experiment was identical to the initial concentration, demonstrating that no denitrification had occurred. When supplemented with molybdate, STM\_ *modA* showed PAO1-like nitrate consumption. Thus, nitrate consumption is molybdate-dependent in the STM\_ *modA* mutant.

It was previously demonstrated that *mod* mutations in *E. coli* caused pleiotropic effects on molybdo enzyme






activity, including nitrate reductase activity [33]. These effects were reversible in the presence of high concentrations of molybdate, which can be internalized by the sulfate transport system under appropriate conditions [16, 34]. However, sulfur compounds have been shown to inhibit the sulfate transport system, thus in protein-rich medium (such as LB), molybdate is likely internalized by another, less specific transporter, presumably the selenite transport system [34]. A novel permease (PerO) internalizing molybdate at micromolar concentrations was identified in *Rhodobacter capsulatus* [35]. PerO also imports sulfate, tungstate and vanadate, suggesting a general oxyanion transporter function. PerO has 30 % AA sequence identity with *P. aeruginosa* PAO1 PA3839, which may be responsible for the nonspecific uptake of molybdate when added at micromolar concentrations, as was done here for the *modA* mutant.

### *modA* and biofilm formation

The *STM\_modA* mutant strain was tested in vitro for biofilm formation, H<sub>2</sub>O<sub>2</sub> sensitivity, heat shock, proteolytic, lipolytic and hemolytic activities, swarming, twitching and swimming motilities, as well as pyocyanin and pyoverdine production. Among all tested phenotypic traits, only biofilm formation presented a statistically significant difference between *STM\_modA* and wild-type PAO1 (Fig. 3a). To confirm whether the weak biofilm formation






observed in *STM\_modA* was caused by a defect in anaerobic growth in the presence of trace amounts of molybdate, the M63 medium was supplemented with molybdate alone, and with molybdate +  $\text{KNO}_3$ . Results showed that the defect was independent from molybdate supplementation, which implies that reduced biofilm formation in *STM\_modA* is not related to its growth defect in anaerobic conditions. When complemented using the recombinant plasmid encoding wild-type ModABC, biofilm formation was only partially restored (Fig. 3b). These results differ from those of Pederick and colleagues [18], according to which biofilm production was unaltered in the *modA* deletion mutant in 5 ml 24 h static cultures. This may be due, at least in part, to important differences between protocols, as biofilm formation is extremely sensitive to growth conditions e.g. [36]. Namely, the significant differences in volume and duration of the assay may have led to different oxygen gradients. If that was the case, it would imply that results from both studies on biofilm formation highlight the complex nature of this process in *P. aeruginosa* and should be interpreted with caution. However, since this study focusses on an insertional mutant, we cannot rule out the possibility of non-specific effects of the mutation, which may also explain this discrepancy. For instance, the gene adjacent to *modA* in the genome of strain PAO1, which was annotated as a putative transcriptional regulator, may be affected by the transposon insertion and result in pleiotropic effects. Further investigation would be required to confirm this gene's function. It is noteworthy, however, that partial phenotype and/or virulence rescue was previously observed in other complemented *P. aeruginosa* mutants see reference [37], namely in mouse models of acute pneumonia and burn sepsis using the same plasmid [38]. Therefore, it is possible that a more general mechanism, which remains to be described, is responsible for these observations.

#### ***modA* is important for virulence during chronic lung infection**

In the original screening experiment where the *STM\_modA* mutant was targeted as defective for in vivo maintenance, pools of 72 *STM* mutant strains were tested simultaneously [15]. Therefore, further investigation was required to precisely describe the involvement of *modA* in the virulence of *P. aeruginosa* PAO1. An in vivo competitive index (CI) in the rat model of chronic lung infection was performed using both wild-type strain PAO1 and the *STM\_modA* mutant. As depicted in Fig. 4, at 7 days post-infection, we recorded an in vivo CI value of 0.004, which represents a 250-fold decrease in mutant colony forming units (CFU) compared to the wild-type strain. Thus, *STM\_modA* is unable to maintain a chronic



lung infection in the rat model when in competition with wild-type PAO1. However, in aerobic combined culture conditions, PAO1 does not outcompete *STM\_modA*, with an in vitro CI value of ~1.0 after 22 h. In addition, *STM\_modA* was capable of normal growth in minimal M9 medium; hence it is not an auxotroph mutant. This contrast between in vitro and in vivo proliferation of the *STM\_modA* mutant strain clearly stresses that mutation of *modA* has consequences that are relatively specific to the context of infection and probably influenced by the interaction with the host.

#### ***modA* is important for virulence in the amoeba model**

In order to further investigate the idea that the role of *modA* in virulence may be affected by interaction with the host, we decided to use a markedly different host model. *D. discoideum* is an alternative model for the study of bacterial virulence where amoebae feed by bacterial uptake using phagocytosis [28, 39]. *P. aeruginosa* has universal virulence factors used to infect phylogenetically diverse hosts, which makes the amoeba model particularly well suited to study its virulence [28, 39]. Previously identified virulence factors using this model, related to quorum sensing and the type III secretion system, were also essential for infection in mammalian and insect models [40]. Many bacterial virulence factors are active against predation by *D. discoideum*, thus virulence is assumed to be inversely proportional to amoeba grazing [29, 39]. In addition, the predation assay is performed in

aerobic conditions where no growth difference has been noticed between PAO1 and the STM\_ *modA* mutant.


As shown in Fig. 5, the STM\_ *modA* mutant was very susceptible to amoeba predation as shown by the presence of phagocytic plaques with only five *D. dictostelium* cells on a bacterial lawn of this mutant compared to PAO1, where a minimum of 5000 *D. dictostelium* cells was required to observe a phagocytic plaque. These results confirmed the virulence defect of the STM\_ *modA* mutant. As *D. discoideum* is an acute infection model while agar-embedded bacteria in the rat lung represent a chronic lung infection model, these two assays highlight the fact that *modA* may be important for the global virulence network of *P. aeruginosa*, even in aerobic conditions.

#### Metabolic and molecular basis of STM\_ *modA* multi-host virulence

Results presented here independently confirmed that transport of the trace anion molybdate is essential for the activity of molybdoenzymes such as nitrate reductase in *P. aeruginosa* [18]. In anaerobic conditions, molybdate uptake was restored in STM\_ *modA* after the addition of molybdate to the culture medium, which suggests less specific internalization of the molecule by a system other than ModABC. The lack of anaerobic growth in STM\_ *modA* was hypothesized to be directly linked to the maintenance defect observed in the rat model of chronic lung infection (CI = 0.004) because this model mimics CF lung infections, which present evidence of oxygen-depletion [7–9, 41, 42]. This hypothesis, however, does not hold in the case of the amoeba predation assay, where a strong virulence defect was observed in at least partly aerobic

conditions; oxygen availability may be limited in a fully grown bacterial lawn. For the biofilm formation assay, our results showed that STM\_ *modA* is unable to produce as much biofilm as PAO1 in vitro, and that this effect is molybdate-independent. As biofilm is a crucial component of *Paeruginosa* chronic lung infections [43], reduced biofilm production in STM\_ *modA* may be related to the virulence defect observed in the rat model of chronic lung infection, in conjunction with anaerobic growth defectiveness. The virulence defect observed in the amoeba model, on the other hand, could be linked to this biofilm defect alone, as for other biofilm-defective mutants shown to have attenuated virulence in the *D. dictostelium* model [44]. Even without consideration for the biofilm results, which have been discussed at length earlier, the amoeba predation assay suggests that the impact of the mutation in *modA* may go beyond its effect on anaerobic growth, perhaps due to pleiotropic effects of this gene.

This study is not the first report of an impact on virulence for a gene implicated in molybdate homeostasis in *P. aeruginosa*. PA1006 encodes a protein of unknown function implicated in molybdate homeostasis that is critical for biofilm maturation and virulence in a lettuce model and in a burned mouse acute virulence model [45]. Other genes implicated in molybdenum utilization and transport (several *moe*, *moa* and *mod* genes) have been linked to a pathogenesis defect in *Mycobacterium tuberculosis* [46]. These genes were required for intracellular growth in macrophages or maintenance in the organs. Interestingly, *modA* has also been identified in a STM experiment with *M. tuberculosis* using a mouse model of acute lung infection with an intravenous administration route [47].



## Conclusions

We have shown that the inactivation of the *modA* gene in a transposon mutant caused a significant defect in *P. aeruginosa* PAO1 for the establishment of chronic lung infection and for resistance to *D. dictostelium* predation. This study complements previous work [18] in providing evidence that molybdate uptake is important for anaerobic growth and multi-host virulence in *P. aeruginosa*. Since the capacity to thrive in anaerobic conditions is relevant to CF lung infections, the *modABC*-encoded transporter may represent a potential target for therapeutic intervention and deserves further investigation.

## Additional file

**Additional file 1: Table S1.** Primers used for sequencing of the insertion site, complementation and sequencing of the vector for complementation.

## Authors' contributions

SP carried out all the experiments except for host model manipulation, performed statistical analyses and drafted the manuscript. JJ performed statistical analyses and drafted the manuscript. IK performed the experiments with the rat model. MMO performed the amoeba predation assay. All authors read and approved final manuscript.

## Author details

<sup>1</sup> Institut de Biologie Intégrative et des Systèmes, Université Laval, Québec, QC, Canada. <sup>2</sup> Département de microbiologie-infectiologie-immunologie, Faculté de Médecine, Université Laval, Québec, QC, Canada. <sup>3</sup> Département de biochimie, de microbiologie et de bio-informatique, Faculté des Sciences et de Génie, Université Laval, Québec, QC, Canada. <sup>4</sup> Centre de recherche de l'Institut universitaire de cardiologie et de pneumologie de Québec (CRI-UCPQ), Québec, QC, Canada.

## Acknowledgements

We are grateful to Anne Sebilo and Manon Couture for support and material used for anaerobic growth condition. We also thank Halim Maaroufi and Brian Boyle for collaboration on this project. The research in R.C.L.'s laboratory is funded by the Canadian Institute for Health Research as part of the CIHR Genomics Program operating grant (MOP-86644) and by Cystic Fibrosis Canada. RCL is a Fonds de la Recherche du Québec - Santé (FRQS) Research Scholar of Exceptional Merit. S.P. is a Master's scholar from CIHR. J.J. is a Cystic Fibrosis Canada postdoctoral fellow. S.J.C. is a research scholar of the FRQS.

## Competing interests

The authors declare that they have no competing interests.

Received: 23 October 2015 Accepted: 6 January 2016

Published online: 12 January 2016

## References

- Rosenfeld M, Gibson RL, McNamara S, Emerson J, Burns JL, Castile R, et al. Early pulmonary infection, inflammation, and clinical outcomes in infants with cystic fibrosis. *Pediatr Pulmonol*. 2001;32(5):356–66. doi:10.1002/ppul.1144.
- Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warren P, Hickey MJ, et al. Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature*. 2000;406(6799):959–64. doi:10.1038/35023079.
- Silby MW, Winstanley C, Godfrey SA, Levy SB, Jackson RW. *Pseudomonas* genomes: diverse and adaptable. *FEMS Microbiol Rev*. 2011;35(4):652–80. doi:10.1111/j.1574-6976.2011.00269.x.
- Balasubramanian D, Schnepfer L, Kumari H, Mathee K. A dynamic and intricate regulatory network determines *Pseudomonas aeruginosa* virulence. *Nucleic Acids Res*. 2013;41(1):1–20. doi:10.1093/nar/gks1039.
- Worlitzsch D, Tarran R, Ulrich M, Schwab U, Cekici A, Meyer KC, et al. Effects of reduced mucus oxygen concentration in airway *Pseudomonas* infections of cystic fibrosis patients. *J Clin Invest*. 2002;109(3):317–25. doi:10.1172/JCI13870.
- Tunney MM, Field TR, Moriarty TF, Patrick S, Doering G, Muhlebach MS, et al. Detection of anaerobic bacteria in high numbers in sputum from patients with cystic fibrosis. *Am J Respir Crit Care Med*. 2008;177(9):995–1001. doi:10.1164/rccm.200708-1151OC.
- Alvarez-Ortega C, Harwood CS. Responses of *Pseudomonas aeruginosa* to low oxygen indicate that growth in the cystic fibrosis lung is by aerobic respiration. *Mol Microbiol*. 2007;65(1):153–65. doi:10.1111/j.1365-2958.2007.05772.x.
- Schober M, Jahn D. Anaerobic physiology of *Pseudomonas aeruginosa* in the cystic fibrosis lung. *Int J Med Microbiol*. 2010;300(8):549–56. doi:10.1016/j.ijmm.2010.08.007.
- Yoon SS, Hennigan RF, Hilliard GM, Ochsner UA, Parvatiyar K, Kamani MC, et al. *Pseudomonas aeruginosa* anaerobic respiration in biofilms: relationships to cystic fibrosis pathogenesis. *Dev Cell*. 2002;3(4):593–603.
- Carlson CA, Ingraham JL. Comparison of denitrification by *Pseudomonas stutzeri*, *Pseudomonas aeruginosa*, and *Paracoccus denitrificans*. *Appl Environ Microbiol*. 1983;45(4):1247–53.
- Arai H. Regulation and Function of Versatile Aerobic and Anaerobic Respiratory Metabolism in *Pseudomonas aeruginosa*. *Front Microbiol*. 2011;2:103. doi:10.3389/fmicb.2011.00103.
- Grunden AM, Shanmugam KT. Molybdate transport and regulation in bacteria. *Arch Microbiol*. 1997;168(5):345–54.
- Frausto da Silva JJR, Williams RJP. The biological chemistry of the elements: the inorganic chemistry of life. Oxford: University press;1993.
- Schwartz G, Hagedoorn P-L, Fischer K. Molybdate and tungstate: uptake, homeostasis, cofactors, and enzymes. In: Nies DH, Silver S, editors. *Molecular Microbiology of Heavy Metals*. Berlin: Springer-Verlag; 2007.
- Potvin E, Lehoux DE, Kukavica-Ibrulj I, Richard KL, Sanschagrin F, Lau GW, et al. *In vivo* functional genomics of *Pseudomonas aeruginosa* for high-throughput screening of new virulence factors and antibacterial targets. *Environ Microbiol*. 2003;5(12):1294–308. doi:10.1046/j.1462-2920.2003.00542.x.
- Rech S, Deppenmeier U, Gunsalus RP. Regulation of the molybdate transport operon, *modABCD*, of *Escherichia coli* in response to molybdate availability. *J Bacteriol*. 1995;177(4):1023–9.
- Self WT, Grunden AM, Hasona A, Shanmugam KT. Molybdate transport. *Res Microbiol*. 2001;152(3–4):311–21.
- Pederick VG, Eijkelkamp BA, Ween MP, Begg SL, Paton JC, McDevitt CA. Acquisition and role of molybdate in *Pseudomonas aeruginosa*. *Appl Environ Microbiol*. 2014;80(21):6843–52. doi:10.1128/aem.02465-14.
- Lehoux DE, Sanschagrin F, Levesque RC. Identification of *in vivo* essential genes from *Pseudomonas aeruginosa* by PCR-based signature-tagged mutagenesis. *FEMS Microbiol Lett*. 2002;210(1):73–80.
- Sanschagrin F, Kukavica-Ibrulj I, Levesque RC. Essential genes in the infection model of *Pseudomonas aeruginosa* PCR-based signature-tagged mutagenesis. *Methods Mol Biol*. 2008;416:61–82. doi:10.1007/978-1-59745-321-9\_5.
- Kukavica-Ibrulj I, Bragonzi A, Paroni M, Winstanley C, Sanschagrin F, O'Toole GA, et al. *In Vivo* Growth of *Pseudomonas aeruginosa* Strains PAO1 and PA14 and the hypervirulent strain LESB58 in a rat model of chronic lung infection. *J Bacteriol*. 2008;190(8):2804–13. doi:10.1128/jb.01572-07.
- Winsor GL, Van Rossum T, Lo R, Khaira B, Whiteside MD, Hancock RE, et al. *Pseudomonas* Genome Database: facilitating user-friendly, comprehensive comparisons of microbial genomes. *Nucleic Acids Res*. 2009;37:483–8. doi:10.1093/nar/gkn861.
- Miranda KM, Espey MG, Wink DA. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide*. 2001;5(1):62–71. doi:10.1006/niox.2000.0319.
- Jeukens J, Boyle B, Kukavica-Ibrulj I, Ouellet MM, Aaron SD, Charette SJ, et al. Comparative genomics of isolates of a *Pseudomonas aeruginosa* epidemic strain associated with chronic lung infections of cystic fibrosis patients. *PLoS One*. 2014;9(2):e87611. doi:10.1371/journal.pone.0087611.
- O'Toole GA. Microtiter dish biofilm formation assay. *J Vis Exp*. 2011;47:2437. doi:10.3791/2437.



26. O'Toole GA, Kolter R. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol.* 1998;28(3):449–61.
27. Cash HA, Woods DE, McCullough B, Johanson WG Jr, Bass JA. A rat model of chronic respiratory infection with *Pseudomonas aeruginosa*. *Am Rev Respir Dis.* 1979;119(3):453–9.
28. Filion G, Charette SJ. Assessing *Pseudomonas aeruginosa* virulence using a nonmammalian host: *Dictyostelium discoideum*. *Method Mol Biol.* 2014;1149:671–80.
29. Froquet R, Lelong E, Marchetti A, Cosson P. *Dictyostelium discoideum*: a model host to measure bacterial virulence. *Nat Protoc.* 2009;4(1):25–30. doi:10.1038/nprot.2008.212.
30. Vizvaryova M, Valkova D. Transposons - the useful genetic tools. *Biologia, Bratislava.* 2004;59(3):309–18.
31. Winsor GL, Lam DK, Fleming L, Lo R, Whiteside MD, Yu NY, et al. *Pseudomonas* genome database: improved comparative analysis and population genomics capability for *Pseudomonas* genomes. *Nucleic Acids Res.* 2011;39:596–600. doi:10.1093/nar/gkq869.
32. Hollenstein K, Dawson RJ, Locher KP. Structure and mechanism of ABC transporter proteins. *Curr Opin Struct Biol.* 2007;17(4):412–8. doi:10.1016/j.sbi.2007.07.003.
33. Glaser JH, DeMoss JA. Phenotypic restoration by molybdate of nitrate reductase activity in *chLD* mutants of *Escherichia coli*. *J Bacteriol.* 1971;108(2):854–60.
34. Lee JH, Wendt JC, Shanmugam KT. Identification of a new gene, *molR*, essential for utilization of molybdate by *Escherichia coli*. *J Bacteriol.* 1990;172(4):2079–87.
35. Gisin J, Muller A, Pfander Y, Leimkuhler S, Narberhaus F, Masepohl B. A *Rhodobacter capsulatus* member of a universal permease family imports molybdate and other oxyanions. *J Bacteriol.* 2010;192(22):5943–52. doi:10.1128/JB.00742-10.
36. O'May CY, Reid DW, Kirov SM. Anaerobic culture conditions favor biofilm-like phenotypes in *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *FEMS Immunol Med Microbiol.* 2006;48(3):373–80.
37. Kukavica-Ibrulj I, Sanschagrin F, Peterson A, Whiteley M, Boyle B, MacKay J, et al. Functional genomics of PycR, a LysR family transcriptional regulator essential for maintenance of *Pseudomonas aeruginosa* in the rat lung. *Microbiology.* 2008;154(7):2106–18. doi:10.1099/mic.0.2007/011239-0.
38. Lau GW, Britigan BE, Hassett DJ. *Pseudomonas aeruginosa* *oxyR* is required for full virulence in rodent and insect models of infection and for resistance to human neutrophils. *Infect Immun.* 2005;73(4):2550–3. doi:10.1128/iai.73.4.2550-2553.2005.
39. Dallaire-Dufresne S, Paquet VE, Charette SJ. *Dictyostelium discoideum*: a model for the study of bacterial virulence. *Can J Microbiol.* 2011;57(9):699–707. doi:10.1139/w11-072.
40. Alibaud L, Kohler T, Coudray A, Prigent-Combaret C, Bergeret E, Perrin J, et al. *Pseudomonas aeruginosa* virulence genes identified in a *Dictyostelium* host model. *Cell Microbiol.* 2008;10(3):729–40. doi:10.1111/j.1462-5822.2007.01080.x.
41. Hassett DJ, Sutton MD, Schurr MJ, Herr AB, Caldwell CC, Matu JO. *Pseudomonas aeruginosa* hypoxic or anaerobic biofilm infections within cystic fibrosis airways. *Trends Microbiol.* 2009;17(3):130–8. doi:10.1016/j.tim.2008.12.003.
42. Palmer KL, Brown SA, Whiteley M. Membrane-bound nitrate reductase is required for anaerobic growth in cystic fibrosis sputum. *J Bacteriol.* 2007;189(12):4449–55. doi:10.1128/JB.00162-07.
43. Hoiby N, Ciofu O, Bjarnsholt T. *Pseudomonas aeruginosa* biofilms in cystic fibrosis. *Future Microbiol.* 2010;5(11):1663–74. doi:10.2217/fmb.10.125.
44. Iwashkiw JA, Seper A, Weber BS, Scott NE, Vinogradov E, Stratilo C, et al. Identification of a general O-linked protein glycosylation system in *Acinetobacter baumannii* and its role in virulence and biofilm formation. *PLoS Pathog.* 2012;8(6):e1002758. doi:10.1371/journal.ppat.1002758.
45. Filiatrault MJ, Tomblin G, Wagner VE, Van Alst N, Rumbaugh K, Sokol P, et al. *Pseudomonas aeruginosa* PA1006, which plays a role in molybdenum homeostasis, is required for nitrate utilization, biofilm formation, and virulence. *PLoS ONE.* 2013;8(2):e55594. doi:10.1371/journal.pone.0055594.
46. Williams M, Mizrahi V, Kana BD. Molybdenum cofactor: a key component of *Mycobacterium tuberculosis* pathogenesis? *Crit Rev Microbiol.* 2013;40:18–29. doi:10.3109/1040841X.2012.749211.
47. Camacho LR, Ensergueix D, Perez E, Gicquel B, Guilhot C. Identification of a virulence gene cluster of *Mycobacterium tuberculosis* by signature-tagged transposon mutagenesis. *Mol Microbiol.* 1999;34(2):257–67. doi:10.1046/j.1365-2958.1999.01593.x.
48. Holloway BW. Genetic recombination in *Pseudomonas aeruginosa*. *J Gen Microbiol.* 1955;13(3):572–81.
49. Schweizer HP. *Escherichia-Pseudomonas* shuttle vector derived from pUC18/19. *Gene.* 1991;97(03838):109–12.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
www.biomedcentral.com/submit

